


Severe Marine Weather Studies using SMOS L-Band Sensor Data and Multi-Sensor Synergies

N. Reul

E. Zabolotskikh, B. Chapron, Y. Quilfen, F. Collard, J. Cotton,
P. Francis, V. Kudryavtsev, J. Tenerelli





❑ Tropical cyclone & Extra-Tropical storm track prediction is steadily improving, while **storm intensity prediction has seen little progress** in the last quarter century.

=> *Important physics are not yet well understood* and implemented in tropical cyclone forecast models.

Missing and unresolved physics, especially at the air-sea interface, are among the factors limiting storm predictions.


❑ Detail Information on **surface winds under Tropical Cyclones** are key to better storm forecasting. However, their **measurements from Space with traditional onboard instruments** (radars, high-frequency radiometers) **is challenging** (rain contamination, lost of sensitivity at very high winds,..)

❑ Focus here: study of low-microwave frequency radiometer capabilities & new inputs from L-band missions (SMOS, SMAP) for ocean surface remote sensing in extreme conditions



Ocean-Atmosphere Interface in very High Wind speed conditions


A breaking wave creates a patch of active foam at its crest – the white cap. As the wave moves on, the leading edge of the white cap follows the breaking crest but the trailing edge remains stationary and is slowly replaced by submerged bubbles in wind-aligned streaks. At very high wind speeds the white cap is blown off the crest in a layer of spray droplets. Under such conditions, the ocean-atmosphere interface is a foam, spray, bubble emulsion layer, which acts as a slip layer for the wind, rather than as a liquid surface [Powell et al., 2003; Emanuel, 2003].




At very high wind speeds this layer covers the waves as a high-velocity white sheet, resulting in white out conditions.



Holthuijsen et al. 2012 investigate these processes using aerial reconnaissance films and GPS drop sondes in hurricanes



Separation of whitecap & streaks coverage



Holthuijsen et al. JGR 2012

Most of the increased surface whitening at & above hurricane force (>33 m/s) is principally induced by the increased streaks coverage- whitecap coverage is found ~constant above Hurricane force ~4 %





Sea Surface Observation Capabilities from Space in Extreme Wind Conditions




Limitations of satellite microwave at high winds


- Active microwave backscatter signal saturates under hurricane force winds and is heavily affected in the presence of high rain rates;
- Contrarily to scatterometer signal, radiometric signal does not saturate with high winds. Moreover, the sensitivity of microwave brightness temperature tends even to increase for the winds above 15 m/s




Donnelly et al. 1998



Rain Anatomy in a hurricane




Rain rate [mm/h]




S.Shen and J. Tenerelli 2007

Wind speed retrieval in extreme winds : SFMR



Increase of the microwave ocean emissivity
with wind speed \leftrightarrow surface foam change impacts



Hurricane hunter P-3




This information can be used to retrieve the surface wind speed in Hurricanes:

Principle of the **Step Frequency Microwave Radiometer (SFMR)** C-band:
=> Use multi-frequency C-band channels to separate wind from rain effects
NOAA's primary airborne sensor for measuring Tropical Cyclone surface wind speeds since 30 year (Ulhorn et al., 2003, 2007).




The Advanced Dvorak Technique (ADT)






The **Advanced Dvorak Technique (ADT)** utilizes longwave-infrared, temperature measurements from geostationary satellites to estimate tropical cyclone (TC) intensity. The ADT is based upon the operational [Dvorak Technique](#) developed by Vern Dvorak of NOAA over 30 years ago..

The **Dvorak Technique** continues to be the standard method for estimating TC intensity where aircraft reconnaissance is not available (all tropical regions outside the North Atlantic and Caribbean Sea), however it has several important limitations and flaws.

SMOS view of Igor Cat 4 Hurricane Igor 2010



Geophysical Model function: $T_b=f(\text{wind speed})$



C-band
L-band:
0.7K/m/s
for hurricanes
0.3 k/(m/s)
below

$$\Delta I = \frac{\Delta(T_H + T_V)}{2} = 0.35 U_{10} - 1.3 \quad U_{10} \leq 33 \text{ m.s}^{-1}$$

$$= 0.75 U_{10} - 14.5 \quad U_{10} \geq 33 \text{ m.s}^{-1}$$

Development of a SMOS wind speed GMF based on Hwind products in IGOR hurricane

Bilinear L-band dependencies with surface wind speed

Reul et al., JGR, 2012



SMOS+STORM Evolution ESA-STSE project


Collaboration IFREMER & Met Office- (2 years: KO Apr 2014)

- Improve high wind speed retrieval algorithms (GMF, rain & wave impacts)
- Produce a Global Tropical Cyclone & Extra-Tropical Cyclone storm catalogue & database from 2010 to now
- Comparisons with NWP models & radiometer & scatterometer data
- Combine with other observations : AMSR2, WindSat, SMAP, CYGNSS
- Evaluate the impact of SMOS High Wind products assimilation on Metoffice forecast
Errors: storm track & intensity forecasts



Detect the useful TC & ETC events in SMOS data: Example of EMILIA

East Pacific TC : EMILIA-2012/07




Position of the Storm center at the time of SMOS Acquisition



Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC :EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07





Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA


East Pacific TC : EMILIA-2012/07






Tasks 2: Detect the useful TC & ETC events in SMOS data: Example of EMILIA

East Pacific TC : EMILIA-2012/07





Monitoring Surface Winds with SMOS in Extra-Tropical Cyclones




SMOS systematically detects higher wind speeds than NWP & could help re-phasing the Storms structures in operational weather forecast models



A view at the SMOS-STORM 2010-2015 TC database

Ensemble of SMOS-TC 320 intercepts considered for Analysis




A subset of 320 SMOS swath intercepts with TCs over 2010-2015, free of Radio Frequency Interferences and with pixel distances >150 km from coasts are selected

Data available at <http://www.smosstorm.org/>




East Pacific SMOS intercepts with 2010-2014 TCs






West Pacific SMOS intercepts with 2010-2014's TCs






South Indian SMOS intercepts with 2010-2014's TCs






North Atlantic SMOS intercepts 2010-2014 TC







Analysing the GMF more in depth and TC intensity meter capability of SMOS

East Pacific TC :JOVA-2011/10



For each TC, multi-incidence Tb contrasts is collected





Storm intensity is evaluated using Best track data and used for further classification




Tropical Storms


Category 1




Category 2



Category 3



Category 4





Saffir–Simpson hurricane wind scale

Category	Wind speeds
Five	≥70 m/s, ≥137 knots ≥157 mph, ≥252 km/h
Four	58–70 m/s, 113–136 knots 130–156 mph, 209–251 km/h
Three	50–58 m/s, 96–112 knots 111–129 mph, 178–208 km/h
Two	43–49 m/s, 83–95 knots 96–110 mph, 154–177 km/h
One	33–42 m/s, 64–82 knots 74–95 mph, 119–153 km/h

Additional classifications



Tropical storm	18–32 m/s, 35–63 knots 39–73 mph, 63–118 km/h
Tropical depression	<17 m/s, <34 knots <38 mph, <62 km/h





Best track eye position

actual eye position

TC eye position is adjusted using 85 GHz datasets




SMOS Tb is recentered on a TC eye-centered frame
and
storm propagation direction is evaluated



SMOS Tb is rotated to a fix North propagation direction for further Tbs averaging


Tropical Storm $34 \leq U_{\max} \leq 64$ knts

119 events intercepted




Category 1 $84 \leq U_{\max} \leq 95$ knts

86 events intercepted



Category 2 : $95 \leq U_{\max} \leq 112$ knts


46 events intercepted



Category 3

$96 \leq U_{\max} \leq 120$ knts


37 events intercepted



Category 4

$113 \leq U_{\max} \leq 135$ knts


27 events intercepted



Category 5


$U_{\max} \geq 135$ knts

(3 events intercepted)











Average L-band Tb contrasts as function of storm Intensity & sectors








Systematic right-hand sectors asymetries in Tb as expected in wind & waves distribution in TCs (extended fetch=>Young, 2003; MacAfee and Bowyer, 2005)



Systematic right-hand sectors asymmetries in Tb as expected in wind & waves distribution in TCs (extended fetch=>Young, 2003; MacAfee and Bowyer, 2005)

When eyewall radii becomes less than instrument resolution (50 km) : eye is not detected






SMOS surface Tb data reveal a clear average growth of amplitude with storm intensity

=> Can be used as a Tropical cyclone intensity meter

SMOS shows sector Distribution asymetries with max in RHS Storm quadrants (east)
=> Information on sea state ?



SMOS STORM SHAKER

New 'average' structural
Information on
tropical cyclones
in terms of radius of high winds

General limits of orbiting scatterometer
Wind speed monitoring capabilities





Derivation of a revised Geophysical Model Function $U=f(Tb)$ using co-located SFMR wind speed data

64 - SFMR flights were co-localized with SMOS-STORM Tb database over 2010-2014


SFMR data from NOAA:

- C-band Tbs
- retrieved surface wind speed (6 km res)
- retrieved rain rate
- SSS along track (climato)
- SST along track (IR data ?)

SMOS data:


- Multi-incidence Tbs
- retrieved wind speed from SMOS 1st GMF
- SST ostia
- SSS from SMOS data composite of L3 during the week preceding each storms

NOAA Hurricane hunter P-3






Derivation of a revised GMF: comparisons with SFMR




Ensemble of SFMR tracks & Wind Speed [knots]




Co-localized SMOS wind Excess Emissivity Δ time (SMOS-SFMR) < 10 hours

Validation: comparison with NOAA/Hurricane Research Division
aircraft data : in moderately strong winds (TS & Cat 1)




North Atlantic TC :leslie-2012/09



SMOS Wind speed -2012/09/07 at -22:19 UTC




Hurricane Leslie 2012/9/7 22:19 UTC




Good agreement



North Atlantic TC : DANIELLE-2010/08




SMOS Wind speed -2010/08/27 at -22:02 UTC






North Atlantic TC : EARL-2010/08




SMOS Wind speed -2010/08/31 at -22:47 UTC






North Atlantic TC : KATIA-2011/09




SMOS Wind speed -2011/09/06 at -09:35 UTC






North Atlantic TC : IRENE-2011/08




SMOS Wind speed -2011/08/25 at -23:13 UTC






North Atlantic TC : TOMAS-2010/11




SMOS Wind speed -2010/11/06 at -22:38 UTC






North Atlantic TC : RAFAEL-2012/10




SMOS Wind speed -2012/10/16 at -09:30 UTC







North Atlantic TC :KARL-2010/09




SMOS Wind speed -2010/09/16 at -11:32 UTC




North Atlantic TC : EARL-2010/09




SMOS Wind speed -2010/09/02 at -23:09 UTC






North Atlantic TC :RINA-2011/10






SMOS Wind speed -2011/10/25 at -11:06 UTC







North Atlantic TC : INGRID-2013/09





Bias!

North Atlantic TC : ISAAC-2012/08



SMOS Wind speed -2012/08/28 at -11:17 UTC
















SMOS+
storms

support to science element






a) EARL: 2010-08-31 @ 22:47 UTC - Vmax = 57.1 m/s








30 co-localisations SMOS/HWIND



New quadratic GMF


$$\Delta I(U_{10}) = SST \cdot (2.7935 \times 10^{-5} U_{10}^2 + 6.8599 \times 10^{-5} U_{10} + 0.0059)$$



SMOS wind RMSE = 5 m/s with small biases above hurricane force
 ECMWF & NCEP RMSE for $u > 33$ m/s = 12-16 m/s



Potential Rain Effects



No clear signal associated with rain but still difficult To firmly conclude




Preparing New Low Microwave frequency merged surface wind Products in TCs

Step1: Merged SMOS+AMSR2 surface winds (on going)

Step2: Merged SMOS+AMSR2+SMAP ...(To be developed in the coming year)

Towards Merged SMOS-AMSR-2-SMAP High wind products




On 18 May 2012 Japan launched a new passive microwave instrument with the largest in the world diameter of antenna - Advanced Microwave Scanning Radiometer (**AMSR2**) onboard Global Change Observation Mission – Water satellite (**GCOM-W1** “Shizuku”)

Additional channel

Better than AMSR-E

AMSR2 Channel Set				
Center Freq. [GHz]	Band width [MHz]	Polarization	Beam width [deg] (Ground res. [km])	Sampling interval [km]
6.925 7.3	350	V & H	<u>1.8 (35 x 61)</u>	10
10.65			<u>1.2 (24 x 41)</u>	
18.7			<u>0.65 (13 x 22)</u>	
23.8			<u>0.75 (15 x 26)</u>	
36.5			<u>0.35 (7 x 12)</u>	
89.0(A&B)	3000		<u>0.15 (3 x 5)</u>	5

Same as AMSR-E



Potential accuracy for SWS retrievals is 1 m/s





An extreme extreme: the super typhoon Hayan in 2013




Super Typhoon
Maximum sustained Wind
Reaching ~150-170 knots

Super Typhoon Haiyan (2013)






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data






Ascat Metop-A & Metop-B Data







Ascat Metop-A & Metop-B Data





Ascat Metop-A & Metop-B Data





Unability of the scatterometer to measure wind speeds above hurricane force (64 knots)

Haiyan Super Typhoon Signature in SMOS data







Figure 1: SMOS retrieved surface wind speed [km/h] along the eye track of super typhoon Haiyan from 4 to 9 Nov 2013.

Haiyan Super Typhoon Signature in SMOS data




Haiyan Typhoon in 2013:
The brightest natural source of L-band radiation ever measured over the oceans
=>an unprecedented natural extreme

Haiyan Super Typhoon Signature in SMOS data




Surface wind speed deduced from the SMOS estimated excess brightness temperature.




Maximum sustained 1 minute wind speed estimated during Haiyan Typhoon. From SMOS data (black filled dots) compared to Advanced Dvorak Technique (ADT=blue diamond), CIMSS (yellow filled dots), SATCON (red) and Best Track from NHC (cyan).

Excellent agreement between SMOS max winds estimates and other traditional Top of the atmosphere estimates datasets (Dvorak, Best track,..)

SMOS versus AMSR2 SWS in Haiyan



Zabolotskikh E.V., L.M. Mitnik, N. Reul, B. Chapron, (2015). New possibilities for geophysical parameter retrievals opened by GCOM-W1 AMSR2. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing (JSTARS), doi: 10.1109/JSTARS.2015.2416514. I F 2.827



Very Coherent L (SMOS) & C (AMSR-2) SWS retrievals 5 hours apart




Figure 20: Top: Superimposed contours of SMOS (dashed) and AMSR2 (filled) surface wind speed fields estimated 5 hours apart as the sensors overpassed the super Typhoon Haiyan on the 7 Nov 2013. Bottom: North-South (left) and East-West (right) sections of the retrieved wind speed through the storm (blue=SMOS; red=AMSR2).



Towards Merged SMOS-AMSR-2-SMAP High wind products









Figure 22: Contours of the merged SMOS+AMSR2 retrieved winds over Haiyan at the threshold levels of 34 (blue), 50 (green) and 64 (orange) knots.



Excellent agreement between SMOS+AMSR2 & Traditional methods of Max wind Estimate (Dvorak)




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/22 at -10:36 UTC




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/22 at -23:08 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/23 at -18:40 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/24 at -07:02 UTC




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/25 at -10:19 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/25 at -18:28 UTC




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/25 at -22:51 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/26 at -06:49 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/26 at -17:32 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/27 at -18:15 UTC




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/27 at -23:13 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/28 at -06:37 UTC




North Atlantic TC : SANDY-2012/10



SMOS Wind speed -2012/10/28 at -10:03 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/28 at -17:20 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/29 at -07:20 UTC




North Atlantic TC : SANDY-2012/10



AMSR2 Wind speed -2012/10/29 at -18:03 UTC

Towards Merged SMOS-AMSR2-SMAP High wind products





Summary

- We evidenced clear SMOS brightness temperature signal (ΔT_B) associated with the passage of Tropical Cyclones
- Correlations between L-band T_b increase with TC intensity from Cat 1 to Cat 5 was demonstrated
- L-band observations provide a first non-atmosphere corrupted view of the ocean surface in extreme conditions \Rightarrow wind speed retrieval with ~ 5 m/s accuracy
- A complete storm database as been generated for the SMOS mission archive:
TC & ETC 2010-now
- We have shown that SMOS can allow to retrieve important structural surface wind features within hurricanes such as the radius of wind speed larger than 34, 50 and 64 knots. These are Key parameters to monitor tropical cyclone intensification
Ascat can provide R34, sometimes R50, but not above R64 \Rightarrow SMOS does



Perspectives

- Merged low-frequency radiometer observations in extremes : SMOS+AMSR-2+SMAP +...CYGNSS=> new opportunity to study air-sea interactions in extreme wind conditions: foam & whitecaps properties, ocean response to TC passage, drag coefficient..
- SMOS wind speed data assimilation experiments into UK Metoffice forecasts model will be performed in the next months to investigate the data impact on:
 - storm track & intensity forecasts skills



TC CELIA - SMOS 2010/06/22 12H18

